

Continuous optimization

Lecture II - Convex optimization

François Glineur

Université catholique de Louvain - EPL/INMA & CORE
Francois.Glineur@uclouvain.be

Inverse Problems and Optimization in Low Frequency Electromagnetism
Continuous and Discrete Optimization workshop March 3rd 2008

Questions and comments ...

... are **more than welcome**, at any time !

Slides **will be** available on the web :

<http://www.core.ucl.ac.be/~glineur/>

References

This lecture's material relies on several references (see at the end), but main ideas can be found in:

- ◇ **Convex Optimization**, Stephen BOYD and Lieven VANDENBERGHE, Cambridge University Press, 2004

Plan for Lecture II - first part

Convex optimization : duality and cones

- ◇ Convex optimization: *definition* and *properties*
- ◇ **Duality** for linear optimization
- ◇ *Duality*: from linear to **conic** optimization

Convex optimization : models and algorithms

- ◇ *Conic* modelling: **three** very expressive cones
- ◇ *Algorithms*: the **interior-point** revolution

Terminology

Possible situations: optimal value

$$\min_{x \in \mathbb{R}^n} f(x) \text{ such that } x \in X \subseteq \mathbb{R}^n$$

Optimal value $f^* = \inf\{f(x) \mid x \in X\}$

- a. $X = \emptyset$: **infeasible** problem (convention: $f^* = +\infty$)
- b. $X \neq \emptyset$: feasible problem ; in this case
 - (a) $f^* > -\infty$: **bounded** problem
 - (b) $f^* = -\infty$: **unbounded** problem

Possible situations: optimal set

$$\min_{x \in \mathbb{R}^n} f(x) \text{ such that } x \in X \subseteq \mathbb{R}^n$$

Optimal value f^* is not always attained

Consider the *optimal set* $X^* = \{x^* \in X \mid f(x^*) = f^*\}$

- a. $X^* \neq \emptyset$: **solvable** problem
(at least one optimal solution)
- b. $X^* = \emptyset$: **unsolvable** problem.

There exists feasible, bounded unsolvable problems !

$$\min \frac{1}{x} \text{ such that } x \in \mathbb{R}_+ \text{ gives } f^* = 0 \text{ but } X^* = \emptyset$$

Convex optimization

Introduction

$$\min f(x) \text{ such that } x \in X$$

A feasible solution x^* is a

◇ **global** minimum iff $f(x^*) \leq f(x) \forall x \in X$

◇ **local** minimum iff there exists an open neighborhood $V(x^*)$ such that

$$f(x^*) \leq f(x) \forall x \in X \cap V .$$

Global minimum \Rightarrow local minimum

Global minima are more interesting but also more difficult to find ... but the notion of **convexity** can help us !

Convexity definitions

◇ A set $S \subseteq \mathbb{R}^n$ is **convex** iff

$$\lambda x + (1 - \lambda)y \in S \quad \forall x, y \in S, \lambda \in [0, 1]$$

◇ A function $f : S \mapsto \mathbb{R}$ is **convex** iff

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \quad \forall x, y, \lambda \in [0, 1]$$

(this imposes that the domain S is convex)

◇ Equivalently, a function $f : S \subseteq \mathbb{R}^n \mapsto \mathbb{R}$ is convex iff its **epigraph** is convex

$$\text{epi } f = \{(x, t) \in \mathbb{R}^n \times \mathbb{R} \mid x \in S \text{ and } f(x) \leq t\}$$

◇ An *optimization* problem is *convex* if it deals with the **minimization** of a convex function on a convex set

Examples

- ◇ $\emptyset, \mathbb{R}^n, \mathbb{R}_+^n, \mathbb{R}_{++}^n$
- ◇ $\{x \mid \|x - a\| < r\}$ and $\{x \mid \|x - a\| \leq r\}$
- ◇ $\{x \mid b^T x < \beta\}$, $\{x \mid b^T x \leq \beta\}$ and $\{x \mid b^T x = \beta\}$
- ◇ In \mathbb{R} : intervals (open/closed, possibly infinite)
- ◇ $x \mapsto c$, $x \mapsto b^T y + \beta_0$, $x \mapsto \|x\|$ and $x \mapsto \|x\|^2$,
 $x \mapsto x^T Q x$ with $Q \in \mathbb{R}^{n \times n}$ positive semidefinite
- ◇ In the case $f : \mathbb{R} \mapsto \mathbb{R}$, we mention $x \mapsto e^x$, $x \mapsto -\log x$, $x \mapsto |x|^p$ with $p \geq 1$.
- ◇ f is **concave** iff $-f$ is convex (i.e. reversing inequalities in the definitions) ; there is no notion of concave set!

Fundamental properties of convex optimization

$$\min_{x \in \mathbb{R}^n} f(x) \text{ such that } x \in X \subseteq \mathbb{R}^n$$

When

- ◇ f is a **convex function** to be **minimized**
- ◇ X is a **convex set**

we are dealing with convex optimization problems and

- ◇ Every **local** minimum is **global**
- ◇ The **optimal set** is **convex**
- ◇ The **KKT** optimality conditions are **sufficient**

Basic properties of convex sets

- ◇ If two sets $S \subseteq \mathbb{R}^n$ and $T \subseteq \mathbb{R}^n$ are convex, so is their **intersection** $S \cap T \subseteq \mathbb{R}^n$
- ◇ If two sets $S \subseteq \mathbb{R}^n$ and $T \subseteq \mathbb{R}^m$ are convex, so is their **Cartesian product** $S \times T \subseteq \mathbb{R}^{n+m}$
- ◇ If f is twice differentiable, we have

$$f \text{ convex} \Leftrightarrow \nabla^2 f \succeq 0$$

- ◇ The only functions that are **simultaneously** convex and concave are the affine functions

Basic properties of convex functions

- ◇ If two functions $f(x)$ and $g(x)$ are convex
 - **Product** $af(x)$ is convex for any scalar $a \geq 0$
 - **Sum** $f(x) + g(x)$ is convex
 - **Maximum** $\max\{f(x), g(x)\}$ is convex

Convexity plays nice with linearity

- ◇ If $S \subseteq \mathbb{R}^n$ is convex and $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^m : x \mapsto Ax + b$ a **linear** function, we have that

$$\Phi S = \{\Phi(x) \mid x \in S\} \text{ is } \mathbf{convex}$$

- ◇ This implies that if $f : x \mapsto f(x)$ is a convex function

$$g : x \mapsto g(x) = f(Ax + b) \text{ is } \mathbf{convex}$$

(but of course **not true** for $af(x) + b$!)

- ◇ Similar result holds for $\Theta : \mathbb{R}^n \rightarrow \mathbb{R}^m : x \mapsto Ax + b$ and

$$\Theta^{-1}S = \{x \mid \Theta(x) \in S\} \text{ is } \mathbf{convex}$$

Feasible set defined with functions

$$X = \{x \in \mathbb{R}^n \mid g_i(x) \leq 0 \text{ and } h_j(x) = 0 \text{ for } i \in I, j \in J\}$$

- ◇ $X_g = \{x \in \mathbb{R}^n \mid g(x) \leq 0\}$ is convex if g is convex
- ◇ When $J = \emptyset$, X is convex when every g_i is convex
- ◇ These two conditions are **not necessary**
- ◇ Allowing now equalities, we note that since $h_i(x) = 0 \Leftrightarrow h_i(x) \leq 0$ and $-h_i(x) \leq 0$, we can guarantee that X is convex when all functions h_i are affine
- ◇ To summarize, X is convex as soon as every g_i is **convex** and **every h_i is affine**

Well-known classes of convex problems

$$\min_{x \in \mathbb{R}^n} f_0(x) \text{ s.t. } f_i(x) \leq 0 \forall i \in I \text{ and } f_i(x) = 0 \forall i \in E$$

- ◇ Linear optimization (LO): f_0 and f_i are **affine** for all $i \in E \cup I$

$$f_i(x) = a_i^T x - b_i$$

- ◇ Quadratically constrained quadratic optimization (QCQO): f_0 and f_i are **convex quadratic** for all $i \in I$

$$f_i(x) = x^T Q_i x + r_i^T x + s_i \text{ with } Q_i \succeq 0$$

(equalities f_i , if present, must still be affine for $i \in E$)

More classes of convex problems

- ◇ Geometric optimization (GO):

f_0 and f_i are **posynomials** (in exponential form)

$$f_i(x) = c_i + \sum_{j \in M_i} \exp(a_{ij}^T x - b_{ij})$$

- ◇ Optimization of p -powers:

f_0 linear, f_i are affine plus sum of convex **powers** with **affine scalar** arguments for all $i \in I$

$$f_i(x) = a_{i0}^T x - b_{i0} + \sum_{j \in M_i} |a_{ij}^T x - b_{ij}|^{p_{ij}} \text{ with } p_{ij} \geq 1$$

Even more classes of convex problems

◇ Sum-of-norm optimization (SNO):

f_0 (and f_i for all $i \in I$, if any) are **convex norms** with affine arguments

$$f_i(x) = \sum_{j \in M_i} \|A_{ij}^T x - B_{ij}\|_{p_{ij}} \quad \text{with } p_{ij} \geq 1$$

$$\text{with } \|y\|_p = \left(|x_1|^p + |x_2|^p + \dots + |x_n|^p\right)^{\frac{1}{p}}$$

◇ Entropy optimization (EO):

f_0 is a sum of **entropy** terms, f_i are affine for all $i \in E$

$$f_0(x) = \sum_i x_i \log x_i \quad (\text{implicitly implying } x \geq 0)$$

◇ Analytic centering (AC):

f_0 is a sum of **logarithmic** terms, f_i are affine for all $i \in I \cup E$

$$f_0(x) = - \sum_{j \in M_0} \log(a_{0j}^T x - b_{0j})$$

Properties of convex optimization

Why is it interesting to consider (or restrict yourself to) convex optimization problems?

Passive features:

- ◇ every local minimum is a **global** minimum
- ◇ set of optimal solutions is **convex**
- ◇ optimality (KKT) conditions are **sufficient** (with regularity assumption)

Any algorithm or solver applied to a convex problem will **automatically** benefit from those features
but there is **more** ...

Properties of convex optimization

Active features:

- ◇ possibility of writing down a **dual** problem strongly related to original problem
(weak duality and, with regularity assumption, strong duality → optimality certificates)
- ◇ possibility of designing dedicated algorithms with **polynomial** algorithmic **complexity**
(in most of the cases: an interior-point method based on the theory of self-concordant barriers)

This is the content of the rest of this lecture

Duality for linear optimization

Standard formulation

Consider the linear problem (with m variables y_i)

$$\max \sum_{i=1}^m b_i y_i \text{ such that } \sum_{i=1}^m a_{ij} y_i \leq c_j \quad \forall 1 \leq j \leq n$$

(objective and n linear inequalities), or

$$\max b^T y \text{ such that } A^T y \leq c$$

(matrix notation with $b, y \in \mathbb{R}^m$, $c \in \mathbb{R}^n$ and $A \in \mathbb{R}^{m \times n}$)

All linear problems can be expressed in this format

When is a problem infeasible ?

In other terms: when is $A^T y \leq c$ **inconsistent** ?

And, more importantly: how can we be **sure** ?

- ◇ Feasible \rightarrow exhibit a feasible solution
- ◇ Infeasible \rightarrow ??

$$3y_1 + 2y_2 \leq 8, \quad -y_2 \leq -3, \quad -y_1 \leq -1$$

Add constraints with weights 1, 2 and 3 to obtain
 $0y_1 + 0y_2 \leq -1 \Leftrightarrow 0 \leq -1 \Leftrightarrow$ a **contradiction**

In general: consider $A^T y \leq c$ or, equivalently, a set of inequalities $a_i^T y \leq c_i$

Proving infeasibility

Multiply each inequality by $a_i^T y \leq c_i$ by a nonnegative constant x_i and take the sum to obtain a **consequence**

$$\sum_{i=1}^n (a_i^T y) x_i \leq \sum_{i=1}^n c_i x_i \text{ with } x_i \geq 0$$

$$\left(\sum_{i=1}^n a_i x_i \right)^T y \leq c^T x \text{ with } x \geq 0$$

$$(Ax)^T y \leq c^T x \text{ with } x \geq 0$$

Contradiction arises only for $0^T y \leq \alpha$ with $\alpha < 0$

This happens **iff** $Ax = 0$ et $c^T x < 0 \rightarrow$ **sufficient** condition for infeasibility but ...

Farkas' Lemma

Theorem: $A^T y \leq c$ is inconsistent if **and only if** there exists $x \geq 0$ such that $Ax = 0$ et $c^T x < 0$

In other words:

Exactly one of the following two systems is consistent

$$Ax = 0, \quad x \geq 0 \quad \text{and} \quad c^T x < 0$$

$$A^T y \leq c$$

Proof relies on topological notions (separation argument)

There always exists a **linear** proof for the infeasibility of a linear system!

Bounds and optimality

Let \bar{y} a feasible solution (satisfying $A^T y \leq c$)
 $\rightarrow b^T \bar{y}$ is a **lower bound** on the optimal value f^*

But how to

- ◇ obtain **upper** bounds on the optimal value ?
- ◇ prove that a feasible solution y^* is optimal ?

Those questions are **linked** since

proving that y^* is optimal
 \Updownarrow
proving that $b^T y^*$ is an upper bound
on the optimal value f^*

Generating upper bounds

Consider

$$\begin{aligned} \max y_1 + 2y_2 + 3y_3 \text{ such that } & y_1 + y_2 \leq 1 & (a) \\ & y_2 + y_3 \leq 2 & (b) \\ & y_3 \leq 3 & (c) \end{aligned}$$

Solution $y = (1, 0, 2)$ is feasible with objective value 7
→ lower bound $f^* \geq 7$

Let us combine constraints: $(a) + (b) + 2(c)$

$$y_1 + y_2 + y_2 + y_3 + 2y_3 \leq 1 + 2 + 2 \times 3 \Leftrightarrow y_1 + 2y_2 + 3y_3 \leq 9$$

→ **upper bound** on the optimal value $f^* \leq 9$

Moreover, considering the feasible solution $y = (2, -1, 3)$ with objective 9 provides a **proof** that $f^* = 9$ is the optimal value of the problem

The best upper bound

Let us find the **best** upper bound using this procedure

$$\max \sum_{i=1}^m b_i y_i \text{ such that } \sum_{i=1}^m a_{ij} y_i \leq c_j \quad \forall 1 \leq j \leq n$$

Introducing again n (multiplying) variables $x_j \geq 0$
we get

$$\sum_{j=1}^n x_j \sum_{i=1}^m a_{ij} y_i \leq \sum_{j=1}^n x_j c_j \Leftrightarrow \sum_{i=1}^m y_i \left(\sum_{j=1}^n a_{ij} x_j \right) \leq \sum_{j=1}^n c_j x_j$$

The best upper bound (continued)

This provides an upper bound on the objective equal to $\sum_{j=1}^n c_j x_j$, assuming that x satisfies

$$\sum_{j=1}^n a_{ij} x_j = b_i \quad \forall 1 \leq i \leq m$$

Minimizing now this upper bound

$$\min \sum_{j=1}^n c_j x_j \quad \text{s.t.} \quad \sum_{j=1}^n a_{ij} x_j = b_i \quad \forall 1 \leq i \leq m \quad \text{and} \quad x_i \geq 0$$

or

$$\min c^T x \quad \text{such that} \quad Ax = b \quad \text{and} \quad x \geq 0$$

We find another **linear optimization** problem which is **dual** to our first problem!

Standard denominations

Using a similar reasoning, we could have started with the **minimization** problem and, looking for the best **lower** bound, derive the original maximization problem

In fact, it is customary in the literature to call

$$\min c^T x \text{ such that } Ax = b \text{ and } x \geq 0$$

the **primal** (P) problem with optimal value p^*
and

$$\max b^T y \text{ such that } A^T y \leq c$$

the **dual** (D) problem with optimal value d^*

Duality properties

- ◇ **Weak duality**: any feasible solution for the primal (resp. dual) provides an upper (resp. lower) bound for the dual (resp. primal)
(immediate consequence of our dualizing procedure)
- ◇ Inequality $b^T y \leq c^T x$ holds for any x, y such that $Ax = b$, $x \geq 0$ and $A^T y \leq c$ (corollary)
- ◇ If the primal (resp. dual) is unbounded, the dual (resp. primal) must be infeasible
(but the converse is **not** true !)

Duality properties (continued)

- ◇ **Strong duality**: If x^* is an optimal solution for the primal, there exists an optimal solution y^* for the dual such that $c^T x^* = b^T y^*$ (in other words: $p^* = d^*$)
- ◇ This property (and its dual) is not trivial, and is a generalization of the Farkas Lemma \rightarrow it is always possible to exhibit a **proof** that a given solution is optimal !
- ◇ However, there are cases where both problems are infeasible: $c = (-1 \ 0)^T$, $b = -1$ et $A = (0 \ 1)$

Other properties and consequences

	$d^* = -\infty$	d^* finite	$d^* = +\infty$
$p^* = -\infty$	Possible, $p^* = d^*$	Impossible	Impossible
p^* finite	Impossible	Possible, $p^* = d^*$	Impossible
$p^* = +\infty$	Possible, $p^* \neq d^*$	Impossible	Possible, $p^* = d^*$

- ◇ One can also write down the dual to a general linear optimization problem
- ◇ One can indifferently solve the primal or the dual to find the optimal objective value
- ◇ **Primal-dual** algorithms solve both problems simultaneously

Conic optimization

Motivation

Objective: **generalize** linear optimization

$$\max b^T y \text{ such that } A^T y \leq c$$

$$\min c^T x \text{ such that } Ax = b \text{ and } x \geq 0$$

while trying to preserve the **nice duality** properties
→ change as little as possible

Idea: generalize the inequalities \leq and \geq

What are properties of **nice** inequalities ?

Generalizing \geq and \leq

Let $K \subseteq \mathbb{R}^n$. Define

$$a \succeq_K 0 \Leftrightarrow a \in K$$

We also have

$$a \succeq_K b \Leftrightarrow a - b \succeq_K 0 \Leftrightarrow a - b \in K$$

as well as

$$a \preceq_K b \Leftrightarrow b \succeq_K a \Leftrightarrow b - a \succeq_K 0 \Leftrightarrow b - a \in K$$

Let us also impose two sensible properties

$$a \succeq_K 0 \Rightarrow \lambda a \succeq_K 0 \quad \forall \lambda \geq 0 \quad (K \text{ is a cone})$$

$$a \succeq_K 0 \text{ and } b \succeq_K 0 \Rightarrow a + b \succeq_K 0$$

(K is closed under addition)

Properties of admissible sets K

- ◇ K is a **convex** set!
- ◇ In fact, if K is a cone, we have

$$K \text{ is closed under addition} \Leftrightarrow K \text{ is convex}$$

Conic optimization

We can then generalize

$$\max b^T y \text{ such that } A^T y \leq c$$

to

$$\max b^T y \text{ such that } A^T y \preceq_K c$$

⇒ This problem is **convex**

The standard linear cases corresponds to $K = \mathbb{R}_+^n$

More requirements for K

◇ $x \succeq 0$ and $x \preceq 0 \Rightarrow x = 0$

which means $K \cap (-K) = \{0\}$ (the cone is **pointed**)

◇ We define the strict inequality by $a \succ 0 \Leftrightarrow a \in \text{int } K$
(and $a \succ b$ iff $a - b \in \text{int } K$)

Hence we require $\text{int } K \neq \emptyset$ (the cone is **solid**)

◇ Finally, we would like to be able to take limits:

If $\{x_i\}_{i \rightarrow \infty}$ with $x_i \succeq_K 0 \forall i$, then $\lim_{i \rightarrow \infty} x_i = \bar{x} \Rightarrow \bar{x} \succeq_K 0$

which is equivalent to saying that K is **closed**

Example: **second-order** (or Lorentz or ice-cream) cone

$$\mathbb{L}^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid \sqrt{x_1^2 + \dots + x_n^2} \leq x_0\}$$

Another example: **semidefinite cone** $K = \mathbb{S}_+^n$ (symmetric positive semidefinite matrices)

Back to conic optimization

A convex cone $K \subseteq \mathbb{R}^n$ that is solid, pointed and closed will be called a **proper** cone

In the following, we will always consider proper cones

We obtain

$$\max_{y \in \mathbb{R}^m} b^T y \text{ such that } A^T y \preceq_K c$$

or, equivalently,

$$\max_{y \in \mathbb{R}^m} b^T y \text{ such that } c - A^T y \in K$$

with problem data $b \in \mathbb{R}^m$, $c \in \mathbb{R}^n$ and $A \in \mathbb{R}^{m \times n}$

Combining several cones

Considering **several conic** constraints

$$A_1^T y \preceq_{K_1} c_1 \text{ and } A_2^T y \preceq_{K_2} c_2$$

which are equivalent to

$$c_1 - A_1^T y \in K_1 \text{ and } c_2 - A_2^T y \in K_2$$

one introduces the **product** cone $K = K_1 \times K_2$ to write

$$(c_1 - A_1^T y, c_2 - A_2^T y) \in K_1 \times K_2$$

$$\Leftrightarrow \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} - \begin{pmatrix} A_1^T \\ A_2^T \end{pmatrix} y \in K_1 \times K_2 \Leftrightarrow \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} - \begin{pmatrix} A_1^T \\ A_2^T \end{pmatrix} y \succeq_{K_1 \times K_2} 0$$

If K_1 and K_2 are proper, $K_1 \times K_2$ is also proper

Equivalence with convex optimization

Conic optimization is clearly a special case of convex optimization: what about the reverse statement ?

$$\min_{x \in \mathbb{R}^n} f(x) \text{ such that } x \in X \subseteq \mathbb{R}^n$$

- ◇ The objective of a convex problem can be assumed **w.l.o.g.** to be **linear** w.l.o.g.: $f(x) = c^T x$
- ◇ The feasible region of a convex problem can be assumed **w.l.o.g.** to be in the **conic** standard format:

$$X = \{x \in K \text{ and } Ax = b\}$$

⇒ conic optimization **equivalent** to convex optimization
Conic format is a **standard form** for convex optimization

A linear objective ?

$$\min_{x \in \mathbb{R}^n} f(x) \text{ such that } x \in X \subseteq \mathbb{R}^n$$



$$\min_{(x,t) \in \mathbb{R}^n \times \mathbb{R}} t \text{ such that } x \in X \text{ and } (x,t) \in \text{epi } f$$



$$\min_{(x,t) \in \mathbb{R}^n \times \mathbb{R}} t \text{ such that } x \in X \text{ and } f(x) \leq t$$

\Rightarrow **equivalent** problem with linear objective

Conic constraints ?

$$K_X = \text{cl}\{(x, u) \in \mathbb{R}^n \times \mathbb{R}_{++} \mid \frac{x}{u} \in X\}$$

is called the (closed) **conic hull** of X

We have that K_X is a **closed convex cone** and

$$x \in X \Leftrightarrow (x, u) \in K_X \text{ and } u = 1$$

$$\min_{x \in \mathbb{R}^n} c^T x \text{ such that } x \in X \subseteq \mathbb{R}^n$$



$$\min_{(x,u) \in \mathbb{R}^n \times \mathbb{R}} c^T x \text{ such that } (x, u) \succeq_{K_X} 0 \text{ and } u = 1$$

\Rightarrow **equivalent** problem with a **conic** constraint

Duality properties

Since we generalized

$$\max b^T y \text{ such that } A^T y \leq c$$

to

$$\max b^T y \text{ such that } A^T y \preceq_K c$$

it is tempting to generalize

$$\min c^T x \text{ such that } Ax = b \text{ and } x \geq 0$$

to

$$\min c^T x \text{ such that } Ax = b \text{ and } x \succeq_K 0$$

But this is **not** the right primal-dual pair !

Dualizing a conic problem

Remembering the dualizing procedure for linear optimization, a **crucial** point lied in the ability to derive consequences by taking **nonnegative linear** combinations of inequalities

Consider now the following statement

$$\begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix} \not\preceq_{\mathbb{L}^2} \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

which is **true** since $(-1)^2 + (-1)^2 \leq 2^2$

Multiplying the first line by 0, 1 and the next two by 1, we get $0.1 \times 2 - 1 \times 1 - 1 \times 1 \geq 0$ or $-1.8 \geq 0$:
 \Rightarrow this is a **contradiction!**

We obtained a contraction although the original system of inequalities was **consistent** \Rightarrow something is wrong!
Some nonnegative linear combinations do not work!

Rescuing duality

Starting with

$$x \in K \subseteq \mathbb{R}^n \Leftrightarrow x \succeq_K 0$$

we identify all vectors (of multipliers) $z \in \mathbb{R}^n$ such that the consequence $z^T x \geq 0$ holds as soon as $x \succeq_K 0$

Hence we define the set

$$K^* = \{z \in \mathbb{R}^n \text{ such that } x^T z \geq 0 \forall x \in K\}$$

The dual cone

$$K^* = \{z \in \mathbb{R}^n \text{ such that } x^T z \geq 0 \ \forall x \in K\}$$

- ◇ For any $x \in K$ and $z \in K^*$, we have $z^T x \geq 0$
- ◇ K^* is a convex cone, called the **dual** cone of K
- ◇ K^* is always **closed**, and if K is closed, $(K^*)^* = K$
- ◇ K is **pointed** (resp. **solid**) $\Rightarrow K^*$ is **solid** (resp. **pointed**)
- ◇ **Cartesian** products: $(K_1 \times K_2)^* = K_1^* \times K_2^*$
- ◇ $(\mathbb{R}_+^n)^* = \mathbb{R}_+^n$, $(\mathbb{L}^n)^* = \mathbb{L}^n$, $(\mathbb{S}_+^n)^* = \mathbb{S}_+^n$:
these cones are **self-dual**
- ◇ But there exists (many) cones that are **not** self-dual

Bounds and optimality

Let \bar{y} a feasible solution (satisfying $A^T y \preceq_K c$)
 $\rightarrow b^T \bar{y}$ is a **lower bound** on the optimal value f^*

But how to

- ◇ obtain **upper** bounds on the optimal value ?
- ◇ prove that a feasible solution y^* is optimal ?

Those questions are **linked** since

proving that y^* is optimal
 \Updownarrow
proving that $b^T y^*$ is an upper bound
on the optimal value f^*

Generating upper bounds

Consider

$$\max 2y_1 + 3y_2 + 2y_3 \text{ such that } \begin{pmatrix} y_1 + y_2 \\ y_2 + y_3 \\ y_3 \end{pmatrix} \preceq_{\mathbb{L}^2} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \begin{matrix} (a) \\ (b) \\ (c) \end{matrix}$$

Solution $y = (-2, 1, 2)$ is **feasible** with objective value 3
→ lower bound $f^* \geq 3$ (since $(2, -1, 1) \in \mathbb{L}^2$)

Let us **combine** constraints: $2(a) + (b) + (c)$
(we have the right to do so since $(2, 1, 1) \in (\mathbb{L}^2)^* = \mathbb{L}^2$)

$$2y_1 + 2y_2 + y_2 + y_3 + y_3 \leq 2 + 2 + 3 \Leftrightarrow 2y_1 + 3y_2 + 2y_3 \leq 7$$

→ **upper bound** on the optimal value $f^* \leq 7$

The best upper bound

Let us find the **best** upper bound using this procedure

$$\max \sum_{i=1}^m b_i y_i \text{ such that } \left(\sum_{i=1}^m a_{ij} y_i \right)_{1 \leq j \leq n} \preceq_K \left(c_j \right)_{1 \leq j \leq n}$$

Introducing again n (multiplying) variables x_j
we get

$$\sum_{j=1}^n x_j \sum_{i=1}^m a_{ij} y_i \leq \sum_{j=1}^n x_j c_j \Leftrightarrow \sum_{i=1}^m y_i \left(\sum_{j=1}^n a_{ij} x_j \right) \leq \sum_{j=1}^n c_j x_j$$

under the **assumption** that $x \in K^*$

The best upper bound (continued)

This provides an upper bound on the objective equal to $\sum_{j=1}^n c_j x_j$, assuming that x satisfies

$$\sum_{j=1}^n a_{ij} x_j = b_i \quad \forall 1 \leq i \leq m$$

Minimizing now this upper bound

$$\min \sum_{j=1}^n c_j x_j \quad \text{s.t.} \quad \sum_{j=1}^n a_{ij} x_j = b_i \quad \forall 1 \leq i \leq m \quad \text{and} \quad x \in K^*$$

or

$$\min c^T x \quad \text{such that} \quad Ax = b \quad \text{and} \quad x \succeq_{K^*} 0$$

We find another **conic optimization** problem which is **dual** to our first problem!

Duality for conic optimization

We have completely mimicked the **dualizing** procedure used for linear optimization

The problem of finding the **best upper bound**

$$\min c^T x \text{ such that } Ax = b \text{ and } x \succeq 0$$

becomes thus

$$\min c^T x \text{ such that } Ax = b \text{ and } x \succeq_{K^*} 0$$

The **correct** primal-dual pair is thus

$$\max b^T y \text{ such that } A^T y \preceq_K c$$

$$\min c^T x \text{ such that } Ax = b \text{ and } x \succeq_{K^*} 0$$

Primal-dual pair

Again, for historical reasons, the min problem is called the primal. Since our cones are closed, $(K^*)^* = K^*$, which means we can write the **primal conic** problem

$$\min c^T x \text{ such that } Ax = b \text{ and } x \succeq_K 0$$

and the **dual conic** problem

$$\max b^T y \text{ such that } A^T y \preceq_{K^*} c$$

- ◇ Very **symmetrical** formulation
- ◇ Computing the dual essentially amounts to **finding K^***
- ◇ All **nonlinearities** are confined to the cones K and K^*

Duality properties

- ◇ **Weak duality**: any feasible solution for the primal (resp. dual) provides an upper (resp. lower) bound for the dual (resp. primal)
(immediate consequence of our dualizing procedure)
- ◇ Inequality $b^T y \leq c^T x$ holds for any x, y such that $Ax = b$, $x \succeq_K 0$ and $A^T y \preceq_{K^*} c$ (corollary)
- ◇ If the primal (resp. dual) is unbounded, the dual (resp. primal) must be infeasible
(but the converse is **not** true!)

Completely similar to the situation for linear optimization

Duality properties (continued)

What about **strong duality** ?

If y^* is an optimal solution for the dual, does there exist an optimal solution x^* for the primal such that $c^T x^* = b^T y^*$ (in other words: $p^* = d^*$) ?

Consider $K = \mathbb{L}^2$ with

$$A = \begin{pmatrix} -1 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, \quad b = (0 \quad -1)^T \quad \text{and} \quad c = (0 \quad 0 \quad 0)^T$$

We can easily check that

- ◇ the primal is **infeasible**
 - ◇ the dual is bounded and **solvable**
- ⇒ strong duality **does not hold** for conic optimization ...

Other troublesome situations

Let $\lambda \in \mathbb{R}_+$: consider

$$\min \lambda x_3 - 2x_4 \text{ s.t. } \begin{pmatrix} x_1 & x_4 & x_5 \\ x_4 & x_2 & x_6 \\ x_5 & x_6 & x_3 \end{pmatrix} \succeq_{\mathbb{S}_+^3} 0, \quad \begin{pmatrix} x_3 + x_4 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

In this case, $p^* = \lambda$ but $d^* = 2$: **duality gap!**

$$\min x_1 \text{ such that } x_3 = 1 \text{ and } \begin{pmatrix} x_1 & x_3 \\ x_3 & x_2 \end{pmatrix} \succeq_{\mathbb{S}_+^2} 0$$

In this case, $p^* = 0$ but the problem is **unsolvable!**

In all cases, one can identify the cause for our troubles: the affine subspace defined by the linear constraints is **tangent** to the cone (it does not intersect its interior)

Rescuing strong duality

A feasible solution to a conic (primal or dual) problem is **strictly** feasible iff it belongs to the **interior** of the cone
In other words, we must have $Ax = b$ and $x \succ_K 0$ for the primal and $A^T y \prec_{K^*} c$ for the dual

Strong duality: If the **dual** problem admits a **strictly** feasible solution, we have either

- ◇ an **unbounded** dual, in which case $d^* = +\infty = p^*$ and the primal is infeasible
- ◇ a **bounded dual**, in which case the primal is **solvable** with $p^* = d^*$ (hence there exists at least one feasible primal solution x^* such that $c^T x^* = p^* = d^*$)

Strong duality (continued)

- ◇ If the **primal** problem admits a **strictly** feasible solution, we have either
 - an **unbounded** primal, in which case $p^* = -\infty = d^*$ and the dual is infeasible
 - a **bounded primal**, in which case the dual is **solvable with $d^* = p^*$** (hence there exists at least one feasible dual solution y^* such that $b^T y^* = d^* = p^*$)
- ◇ The first case is a mere consequence of weak duality
- ◇ Finally, when both problems admit a strictly feasible solution, both problems are **solvable** and we have

$$c^T x^* = p^* = d^* = b^T y^*$$

Conic modelling with three cones

A first cone: \mathbb{R}_+^n

Standard meaning for inequalities:

$$\succ_{\mathbb{R}_+^n} \Leftrightarrow \geq$$

\Rightarrow linear optimization

But we can also model some nonlinearities!

$$|x_1 - x_2| \leq 1 \Leftrightarrow -1 \leq x_1 - x_2 \leq 1$$

$$|x_1 - x_2| \leq t \Leftrightarrow \begin{pmatrix} x_1 - x_2 - t \\ x_2 - x_1 - t \end{pmatrix} \leq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Terminology: conic representability

- ◇ Set S is K -representable if can be expressed as feasible region of conic problem using cone K
- ◇ Closed under intersection and Cartesian product
- ◇ Function f is K -representable iff its epigraph is K -representable
- ◇ Closed under sum, positive multiplication and max
- ◇ What we can do in practice: minimize a K -representable function over a K -representable set where K is a product of cones \mathbb{R}_+^n , \mathbb{L}^n , \mathbb{S}_+^n and \mathbb{R}^n

A simple example

Consider set

$$S = \{x_1^2 + x_2^2 \leq 1\}$$

→ can be modelled as

$$(x_0, x_1, x_2) \in \mathbb{L}^2 \text{ and } x_0 = 1$$

⇒ S is \mathbb{L}^2 -representable

but an additional variable x_0 was needed

⇒ formally, $S \subseteq \mathbb{R}^n$ is K -representable

iff there exists a set $T \subseteq \mathbb{R}^{n+m}$ such that

a. T is K -representable

b. $x \in S$ iff there exists $t \in \mathbb{R}^m$ such that $(x, t) \in T$

(i.e. S is the projection of T on \mathbb{R}^n)

Back to \mathbb{R}_+^n

- ◇ Polyhedrons and polytopes are \mathbb{R}_+^n -representable
- ◇ Hyperplanes and half-planes are \mathbb{R}_+^n -representable
- ◇ Affine functions $x \mapsto a^T x + b$ are \mathbb{R}_+^n -representable
- ◇ Absolute values $x \mapsto |a^T x + b|$ are \mathbb{R}_+^n -representable
- ◇ *Convex* piecewise linear functions are \mathbb{R}_+^n -representable

Two potential **issues** with \mathbb{R}_+^n :

a. free variables in the primal $\rightarrow x = x^+ - x^-$

b. equalities in the dual $\rightarrow a^T x \leq c$ and $a^T x \geq c$

But these are **wrong** solutions !

What use is $K = \mathbb{R}^n$?

◇ $K = \mathbb{R}^n$ and $K^* = \{0\}$

◇ Can be used to introduce **free** variables in the primal
 $Ax = b, x \succeq_K 0$

$$x \succeq_{\mathbb{R}^n} 0 \quad \Leftrightarrow \quad x \text{ is free}$$

◇ or **equalities** in the dual $A^T y \preceq_{K^*} c$

$$A^T y \preceq_{\{0\}} c \quad \Leftrightarrow \quad A^T y = c$$

in combination with other cones

◇ \mathbb{R}^n in dual or $\{0\}$ in primal is useless!

What use is \mathbb{L}^n ?

$$\diamond f : x \mapsto \|x\|, f : x \mapsto \|x\|^2 \text{ and } f : (x, z) \mapsto \frac{\|x\|^2}{z}$$

$$\diamond B_r = \{x \in \mathbb{R}^n \mid \|x\| \leq r\}$$

$$\diamond \{(x, y) \in \mathbb{R}_+^2 \mid xy \geq 1\}$$

$$\diamond \{(x, y, z) \in \mathbb{R}_+^2 \times \mathbb{R} \mid xy \geq z^2\}$$

$$\diamond \{(a, b, c, d) \in \mathbb{R}_+^4 \mid abcd \geq 1\}$$

$$\diamond \{(x, t) \in \mathbb{R}^n \times \mathbb{R} \mid x^T Q x \leq t\} \text{ with } Q \in \mathbb{S}_+^n$$

\Rightarrow **second-order** cone optimization

Very useful trick: $xy \geq z^2 \Leftrightarrow (x + y, x - y, 2z) \in \mathbb{L}^2$

Unfortunately, $(x, y) \mapsto \frac{x}{y}$ is **not** convex!

What use is \mathbb{S}_+^n ?

Preliminary remark: for the purpose of conic optimization, members of \mathbb{S}^n are viewed as **vectors** in $\mathbb{R}^{n \times n}$

What about **constraint** $Ax = b$?

$$Ax = b \Leftrightarrow a_i^T x = b_i \quad \forall i$$

$a_i^T x$ can be viewed as the inner product between a_i and x

Let $X, Y \in \mathbb{S}^n$: their **inner product** is

$$X \bullet Y = \sum_{1 \leq i, j \leq n} X_{i,j} Y_{i,j} = \text{trace}(XY)$$

→ replace $a_i^T x$ by $A_i \bullet X$ with $A_i, X \in \mathbb{S}^n$

Standard format for semidefinite optimization

The **primal** becomes

$\min C \bullet X$ such that $A_i \bullet X = b_i \forall 1 \leq i \leq m$ and $X \succeq 0$

In the conic dual, we have

$$A^T y = \sum a_i y_i, \text{ an application from } \mathbb{R}^m \mapsto \mathbb{R}^n$$

\Rightarrow with the \mathbb{S}_+^n cone, we have

$$\mathcal{A}(y) = \sum A_i y_i, \text{ an application from } \mathbb{R}^m \mapsto \mathbb{S}^n$$

which gives for the **dual**

$$\max b^T y \text{ such that } \sum_{i=1}^m A_i y_i \preceq C$$

What use is \mathbb{S}_+^n (continued) ?

- ◇ \mathbb{S}_+^n generalizes both \mathbb{R}_+^n and \mathbb{L}^n (arrow matrices)
(however, using \mathbb{R}_+^n and \mathbb{L}^n is more efficient)
- ◇ $f : X \mapsto \lambda_{max}(X)$ and $f : X \mapsto -\lambda_{min}(X)$
- ◇ $f : X \mapsto \max_i |\lambda_i| (X)$ (spectral norm)
- ◇ Describing ellipsoids $\{x \in \mathbb{R}^n \mid (x-c)^T E (x-c) \leq 1\}$
with $E \succeq 0$
- ◇ Matrix constraint $XX^T \preceq Y$
using the **Schur Complement** lemma
When $A \succ 0$: $\begin{pmatrix} A & B \\ B^T & C \end{pmatrix} \succeq 0 \Leftrightarrow C - B^T A^{-1} B \succeq 0$
- ◇ And more ...

Interior-point methods

Back to convex optimization

Let $f : \mathbb{R}^n \mapsto \mathbb{R}$ be a convex function, $C \subseteq \mathbb{R}^n$ be a convex set : optimize a vector $x \in \mathbb{R}^n$

$$\inf_{x \in \mathbb{R}^n} f(x) \quad \text{s.t.} \quad x \in C \quad (\text{P})$$

Properties

- ◇ All local optima are *global*, optimal set is **convex**
- ◇ Lagrange duality \rightarrow **strongly related** dual problem
- ◇ Objective can be taken linear **w.l.o.g.** ($f(x) = c^T x$)

Principle

Approximate a constrained problem by

a *family* of **unconstrained** problems

Use a **barrier** function F to replace the inclusion $x \in C$


- ◇ F is smooth
- ◇ F is strictly convex on $\text{int } C$
- ◇ $F(x) \rightarrow +\infty$ when $x \rightarrow \partial C$

$$\rightarrow C = \text{cl dom } F = \text{cl } \{x \in \mathbb{R}^n \mid F(x) < +\infty\}$$

Central path

Let $\mu \in \mathbb{R}_{++}$ be a parameter and consider

$$\inf_{x \in \mathbb{R}^n} \frac{c^T x}{\mu} + F(x) \quad (\mathbf{P}_\mu)$$



$$x_\mu^* \rightarrow x^* \text{ when } \mu \searrow 0$$

where

- ◇ x_μ^* is the (unique) solution of (\mathbf{P}_μ) (\rightarrow central path)
- ◇ x^* is a solution of the original problem (\mathbf{P})

Ingredients

- ◇ A method for **unconstrained** optimization
- ◇ A barrier function

Interior-point methods rely on

- ◇ **Newton's method** to compute x_μ^*
- ◇ When C is defined with convex constraints $g_i(x) \leq 0$, one can introduce the **logarithmic** barrier function

$$F(x) = - \sum_{i=1}^n \log(-g_i(x))$$

Question: What is a good barrier, i.e. a barrier for which Newton's method is efficient ?

Answer: A *self-concordant* barrier

Self-concordant barriers

Definition [Nesterov & Nemirovski, 1988]

$F : \text{int } C \mapsto \mathbb{R}$ is called (κ, ν) -self-concordant on C iff

- ◇ F is convex
- ◇ F is three times differentiable
- ◇ $F(x) \rightarrow +\infty$ when $x \rightarrow \partial C$
- ◇ the following **two** conditions hold

$$\nabla^3 F(x)[h, h, h] \leq 2\kappa \left(\nabla^2 F(x)[h, h] \right)^{\frac{3}{2}}$$

$$\nabla F(x)^\top (\nabla^2 F(x))^{-1} \nabla F(x) \leq \nu$$

for all $x \in \text{int } C$ and $h \in \mathbb{R}^n$

A (simple?) example

For **linear** optimization, $C = \mathbb{R}_+^n$: take $F(x) = -\sum_{i=1}^n \log x_i$

When $n = 1$, we can choose $(\kappa, \nu) = (1, 1)$

$$\diamond \nabla F(x) = -\frac{1}{x} \text{ and } \nabla F(x)^\top h = -\frac{h}{x}$$

$$\diamond \nabla^2 F(x) = \frac{1}{x^2} \text{ and } \nabla^2 F(x)[h, h] = \frac{h^2}{x^2}$$

$$\diamond \nabla^3 F(x) = -2\frac{1}{x^3} \text{ and } \nabla^3 F(x)[h, h, h] = -2\frac{h^3}{x^3}$$

When $n > 1$, we have

$$\diamond \nabla F(x) = (-x_i^{-1}) \text{ and } \nabla F(x)^\top h = -\sum h_i x_i^{-1}$$

$$\diamond \nabla^2 F(x) = \text{diag}(x_i^{-2}) \text{ and } \nabla^2 F(x)[h, h] = \sum h_i^2 x_i^{-2}$$

$$\diamond \nabla^3 F(x) = \text{diag}_3(-2x_i^{-3}), \nabla^3 F(x)[h, h, h] = -2 \sum h_i^3 x_i^{-3}$$

and one can show that $(\kappa, \nu) = (1, n)$ is valid

Barrier calculus

Two elementary results:

◇ **Scaling:**

F is a (κ, ν) -s.-c. barrier for $\mathcal{C} \subseteq \mathbb{R}^n$ and $\lambda \in \mathbb{R}_{++}$
 $\Rightarrow (\lambda F)$ is a $(\frac{\kappa}{\sqrt{\lambda}}, \lambda\nu)$ -s.-c. barrier for \mathcal{C}

◇ **Sum:**

F is a (κ_1, ν_1) -s.-c. barrier for $\mathcal{C}_1 \subseteq \mathbb{R}^n$

G is a (κ_2, ν_2) -s.-c. barrier for $\mathcal{C}_2 \subseteq \mathbb{R}^n$

$\Rightarrow (F + G)$ is a $(\max\{\kappa_1, \kappa_2\}, \nu_1 + \nu_2)$ -s.-c. barrier
for the set $\mathcal{C}_1 \cap \mathcal{C}_2$ (if nonempty)

Complexity result

Summary

Self-concordant barrier \Rightarrow polynomial number of iterations to solve (P) within a given accuracy

Short-step method: follow the central path

- ◇ **Measure** distance to the central path with $\delta(x, \mu)$
- ◇ Choose a starting iterate with a **small** $\delta(x_0, \mu_0) < \tau$
- ◇ While accuracy is not attained
 - a. Decrease μ geometrically (δ **increases** above τ)
 - b. Take a Newton step to minimize barrier (δ **decreases** below τ)

Geometric interpretation

Two self-concordancy conditions: each has its role

- ◇ Second condition bounds the size of the Newton step
⇒ **controls** the **increase** of the distance to the central path when μ is updated
- ◇ First condition bounds the variation of the Hessian
⇒ guarantees that the Newton step **restores** the initial **distance** to the central path

Summarized complexity result

$$\mathcal{O} \left(\kappa \sqrt{\nu} \log \frac{1}{\epsilon} \right)$$

iterations lead a solution with **ϵ accuracy** on the **objective**

Complexity result

- ◇ Let F be a (κ, ν) -self-concordant barrier for C and let $x_0 \in \text{int } C$ be a starting point, a **short-step interior-point** algorithm can solve problem (P) up to ϵ accuracy within

$$\mathcal{O} \left(\kappa \sqrt{\nu} \log \frac{c^T x_0 - p^*}{\epsilon} \right) \text{ iterations,}$$

such that at each iteration the self-concordant barrier and its first and second derivatives have to be evaluated and a linear system has to be solved in \mathbb{R}^n

- ◇ Complexity **invariant** w.r.t. to **scaling** of F
- ◇ Universal bound on complexity parameter: $\kappa \sqrt{\nu} \geq 1$

Corollary

Assume F , ∇F and $\nabla^2 F$ are **polynomially** computable
 \Rightarrow problem (P) can be solved in **polynomial** time

Existence

There exists a **universal** SC barrier with parameters

$$\kappa = 1 \text{ and } \nu = \mathcal{O}(n)$$

(**but** not necessarily efficiently computable)

Examples

- ◇ linear optimization: $(\kappa, \nu) = (1, n) \Rightarrow \mathcal{O}(\sqrt{n} \log \frac{1}{\varepsilon})$
- ◇ entropy optimization: $\kappa = 1$ and $\nu = 2n \Rightarrow \mathcal{O}(\sqrt{n} \log \frac{1}{\varepsilon})$
($\inf c^T x + \sum_i x_i \log x_i$ such that $Ax = b$ and $x \geq 0$)

Sketch of the proof

Define $n_\mu(x)$ the **Newton step** taken from x to x_μ^*

$$n_\mu(x) = 0 \text{ if and only if } x = x_\mu^*$$

We take

$$\delta(x, \mu) = \|n_\mu(x)\|_x \quad (\text{size of the Newton step})$$

with a well-chosen (*coordinate invariant*) norm $\|\cdot\|_x$

Set $k \leftarrow 0$, perform the following **main loop**:

a. $\mu_{k+1} \leftarrow \mu_k(1 - \theta)$ (*decrease barrier param*)

b. $x_{k+1} \leftarrow x_k + n_{\mu_{k+1}}(x_k)$ (*take Newton step*)

c. $k \leftarrow k + 1$

Sketch of the proof (continued)

Key choice: parameters τ and θ such that

$$\delta(x_k, \mu_k) < \tau \quad \Rightarrow \quad \delta(x_{k+1}, \mu_{k+1}) < \tau$$

To relate $\delta(x_k, \mu_k)$ and $\delta(x_{k+1}, \mu_{k+1})$,
introduce an **intermediate** quantity

$$\delta(x_k, \mu_{k+1})$$

We will also denote for simplicity

$$x_k \leftrightarrow x$$

$$\mu_k \leftrightarrow \mu$$

Sketch of the proof (end)

Given a (κ, ν) -self-concordant barrier:

◇ $x \in \text{dom } F$ and $\mu^+ = (1 - \theta)\mu \Rightarrow$

$$\delta(x, \mu^+) \leq \frac{\delta(x, \mu) + \theta\sqrt{\nu}}{1 - \theta}$$

◇ $x \in \text{dom } F$ and $\delta(x, \mu) < \frac{1}{\kappa} \Rightarrow$ define $x^+ = x + n_\mu(x)$

$$x^+ \in \text{dom } F \text{ and } \delta(x^+, \mu) \leq \kappa \left(\frac{\delta(x, \mu)}{1 - \kappa\delta(x, \mu)} \right)^2$$

with e.g. possible choice for parameters

$$\tau = \frac{1}{4\kappa} \text{ and } \theta = \frac{1}{16\kappa\sqrt{\nu}}$$

(hence the name short-step)

Primal-dual algorithms

Advantage of **conic** optimization over **standard** convex optimization is (symmetric) **duality**

However previous approach does **not** seem to use it !
 \Rightarrow a **better** approach that uses duality is needed

The linear case (again)

Introduce additional vector of variables $s \in \mathbb{R}^n$

$$\min c^T x \text{ such that } Ax = b \text{ and } x \geq 0$$

and

$$\max b^T y \text{ such that } A^T y + s = c \text{ and } s \geq 0$$

Primal-dual optimality conditions

$$\min c^T x \text{ such that } Ax = b \text{ and } x \geq 0$$

and

$$\max b^T y \text{ such that } A^T y + s = c \text{ and } s \geq 0$$

Duality tells us x^* and y^* are **optimal iff** they satisfy

$$Ax = b, x \geq 0, A^T y + s = c, s \geq 0 \text{ and } c^T x = b^T y$$

or

$$Ax = b, x \geq 0, A^T y + s = c, s \geq 0 \text{ and } x_i s_i = 0 \forall i$$

Both problems are handled **simultaneously**

Perturbed optimality conditions

Introducing a **logarithmic barrier** term in both problems

$$\min c^T x - \mu \sum_i \log x_i \text{ such that } Ax = b \text{ and } x > 0$$

$$\max b^T y + \mu \sum_i \log s_i \text{ such that } A^T y + s = c \text{ and } s > 0$$

one can derive new **perturbed** optimality conditions

$$Ax = b, x \geq 0, A^T y + s = c, s \geq 0 \text{ and } x_i s_i = \mu \forall i$$

Again, **both** problems are handled **simultaneously**

Primal-dual path following algorithm

Same principle as in the general case:

- ◇ Follow the **central** path
- ◇ **Not** wandering **too far** from it
- ◇ Until (primal-dual) **optimality**
- ◇ Using a **polynomial** number of iterations

Complexity is also the same:

$$\mathcal{O} \left(\sqrt{n} \log \frac{1}{\varepsilon} \right) \text{ iterations to get } \varepsilon \text{ accuracy}$$

But this scheme is **very efficient in practice** (long steps)
(all practical implementations use it nowadays)

What about other convex/conic problems ?

This **primal-dual** scheme is only generalizable to cones that are

a. **self-dual** ($K = K^*$)

b. **homogeneous**

(linear automorphism group acts transitively on $\text{int } K$)

(*[Nesterov & Todd 97]*)

There exists a **complete classification** of these cones :
in the real case, they are ...

$$\mathbb{R}_+^n, \quad \mathbb{L}^n \quad \text{and} \quad \mathbb{S}_+^n$$

and their Cartesian products!

Complexity

Complexity for a product of $\mathbb{R}_+^n, \mathbb{L}^n, \mathbb{S}_+^n$

$$\mathcal{O}\left(\sqrt{\nu} \log \frac{1}{\varepsilon}\right) \text{ iterations to get } \varepsilon \text{ accuracy}$$

where ν is the sum of

◇ n for \mathbb{R}_+^n (see above) (barrier term is $-\sum \log x_i$)

◇ n for \mathbb{S}_+^n (although there are $n(n+1)/2$ variables)
(barrier term is $-\log \det X = -\sum \log \lambda_i$)

◇ 2 for \mathbb{L}^n (independently of n !)

(barrier term is $-\log(x_0^2 - \sum x_i^2)$; no $-\log x_0$ term!)

→ these problems are solved **very efficiently in practice**

More applications

Using **semidefinite** optimization:

Positive polynomials

- ◇ Single variable case: **exact** formulation
 - ◇ **Test** positivity and **minimize** on an interval
 - ◇ Multiple variable case: **relaxation** only
-

The MAX-CUT relaxation

- ◇ **Relaxation** of a difficult discrete problem
- ◇ With a quality **guarantee** (0.878)

References

Convex optimization

- ◇ Convex Analysis, ROCKAFELLAR, Princeton University Press, 1980
- ◇ Convex optimization, BOYD and VANDENBERGHE, Cambridge University Press, 2004 (on the web)

Convex modelling

- ◇ Lectures on Modern Convex Optimization, Analysis, Algorithms, and Engineering Applications, BEN-TAL and NEMIROVSKI, MPS/SIAM Series on Optimization, 2001

Interior-point methods (linear)

- ◇ Primal-Dual Interior-Point Methods, WRIGHT
SIAM, 1997
- ◇ Theory and Algorithms for Linear Optimization, ROOS,
TERLAKY, VIAL, John Wiley & Sons, 1997

Interior-point methods (convex)

- ◇ Interior-point polynomial algorithms in convex programming, NESTEROV & NEMIROVSKI, SIAM, 1994
- ◇ A Mathematical View of Interior-Point Methods in Convex Optimization, RENEGAR,
MPS/SIAM Series on Optimization, 2001

Semidefinite optimization applications

- ◇ Handbook of Semidefinite Programming,
WOLKOWICZ, SAIGAL, VANDENBERGHE (eds.)
Kluwer, 2000
- ◇ Semidefinite programming, BOYD, VANDENBERGHE,
SIAM Review 38 (1), 1996

Software: two choices among many others

- ◇ Linear & second-order cone: **MOSEK** (commercial)
- ◇ Linear, sec.-ord. & semidefinite: **SeDuMi** (free)

Thanks for you attention

Does linear optimization exist at all ?

Let us only mention the following *not so well-known* theorem, due to Dr. **Addock PRILFIRST**

Theorem

The objective function of any linear program is **constant** on its feasible region

Proof

$$\begin{aligned} & \{ \min c^T x \mid Ax = b, x \geq 0 \} = \{ \max b^T y \mid A^T y \leq c \} \\ & \geq \{ \min b^T y \mid A^T y \leq c \} = \{ \max c^T x \mid Ax = b, x \leq 0 \} \\ & \geq \{ \min c^T x \mid Ax = b, x \leq 0 \} = \{ \max b^T y \mid A^T y \geq c \} \\ & \geq \{ \min b^T y \mid A^T y \geq c \} = \{ \max c^T x \mid Ax = b, x \geq 0 \} \\ & \geq \{ \min c^T x \mid Ax = b, x \geq 0 \} \end{aligned}$$

Thanks for you attention